

COHOMOLOGY WITH LOCAL COEFFICIENTS OF SOLVMANIFOLDS AND MORSE-NOVIKOV THEORY

DMITRI V. MILLIONSCHIKOV

ABSTRACT. We study the cohomology $H_{\lambda\omega}^*(G/\Gamma, \mathbb{C})$ of the deRham complex $\Lambda^*(G/\Gamma) \otimes \mathbb{C}$ of a compact solvmanifold G/Γ with a deformed differential $d_{\lambda\omega} = d + \lambda\omega$, where ω is a closed 1-form. This cohomology naturally arises in the Morse-Novikov theory. We show that for a solvable Lie group G with a completely solvable Lie algebra $\mathfrak{g}$ and a cocompact lattice $\Gamma \subset G$ the cohomology $H_{\lambda\omega}^*(G/\Gamma, \mathbb{C})$ coincides with the cohomology $H_{\lambda\omega}^*(\mathfrak{g})$ of the Lie algebra $\mathfrak{g}$ associated with the one-dimensional representation $\rho_{\lambda\omega} : \mathfrak{g} \rightarrow \mathbb{K}, \rho_{\lambda\omega}(\xi) = \lambda\omega(\xi)$. Moreover $H_{\lambda\omega}^*(G/\Gamma, \mathbb{C})$ is non-trivial if and only if $-\lambda[\omega]$ belongs to the finite subset $\{0\} \cup \bar{\Omega}_{\mathfrak{g}}$ in $H^1(G/\Gamma, \mathbb{C})$ well defined in terms of $\mathfrak{g}$.

INTRODUCTION

In the begining of the 80-th S.P. Novikov constructed ([N1], [N2]) an analogue of the Morse theory for smooth closed 1-forms on a compact smooth manifold M . In particular he introduced the Morse-type inequalities (Novikov's inequalities) for numbers $m_p(\omega)$ of zeros of index p of a closed 1-form ω on M . A lot of papers was devoted to this problem in the following years (see [N4] for references). In [N3], [Pa] a method of obtaining the torsion-free Novikov inequalities in terms of the deRham complex of manifold was proposed. This method was based on Witten's approach [W] to the Morse theory. A. Pazhitnov obtained some important results in this direction in [Pa]. The cohomology of the deRham complex $\Lambda^*(M)$ with the deformed differential $d + \lambda\omega$ coincides with the cohomology $H_{\rho_{\lambda\omega}}^*(M, \mathbb{C})$ with coefficients in the local system $\rho_{\lambda\omega}$ of groups $\mathbb{C}$, $\rho_{\lambda\omega}(\gamma) = \exp \int_{\gamma} \lambda\omega$ and for sufficiently large real numbers λ we have the following estimate (see [Pa]):

$$m_p(\omega) \geq \dim H_{\rho_{\lambda\omega}}^p(M, \mathbb{C}), \forall p.$$

L. Alania in [Al] studied $H_{\rho_{\lambda\omega}}^*(M_n, \mathbb{C})$ of a class of nilmanifolds M_n . He proved that $H_{\rho_{\lambda\omega}}^*(M_n, \mathbb{C})$ is trivial if $\lambda\omega \neq 0$. The partial answer for the case $\lambda\omega = 0$ was obtained by the present author in [Mill]. In both cases the proof was based on the Nomizu theorem [Nz] and the computations were made in terms of the corresponding nilpotent Lie algebra $\mathcal{V}_n$. The starting point of this article was the intention to improve the results of [Al] in more general situation considering solvmanifolds and to find examples of manifolds M with non-trivial $H_{\rho_{\lambda\omega}}^*(M, \mathbb{C})$, $\lambda\omega \neq 0$. One of the first observations that was made in this direction: *for a nilmanifold G/Γ the cohomology $H_{\lambda\omega}^*(G/\Gamma, \mathbb{C})$ coincides with the cohomology $H_{\lambda\omega}^*(\mathfrak{g})$ associated with the*

1991 *Mathematics Subject Classification.* 58A12, 17B30, 17B56 (Primary) 57T15 (Secondary).

Key words and phrases. Solvmanifolds, nilmanifolds, cohomology, local system, Morse-Novikov theory, solvable Lie algebras .

Partially supported by the Russian Foundation for Fundamental Research, grant no. 99-01-00090 and PAI-RUSSIE, dossier no. 04495UL.

one-dimensional representation of the Lie algebra $\rho_{\lambda\omega} : \mathfrak{g} \rightarrow \mathbb{C}, \rho_{\lambda\omega}(\xi) = \lambda\omega(\xi)$ and hence $H_{\lambda\omega}^*(\mathfrak{g}) = 0$ by Dixmier's theorem [D](Corollary 2.5).

Applying Hattori's theorem [H] one can observe that the isomorphism $H_{\lambda\omega}^*(G/\Gamma, \mathbb{C}) \cong H_{\lambda\omega}^*(\mathfrak{g})$ still holds on for compact solvmanifolds G/Γ with completely solvable Lie group G . A kind of minimal model of a solvable Lie algebra $\mathfrak{g}$, the free d -algebra $(\Lambda^*(\omega_1, \dots, \omega_n), d)$ with differential d

$$d\omega_i = 0, i = 1, \dots, k; \quad d\omega_j = \alpha_j \wedge \omega_j + P_j(\omega_1, \dots, \omega_{j-1}), j = k+1, \dots, n.$$

is considered (Lemma 3.2). By means of $(\Lambda^*(\omega_1, \dots, \omega_n), d)$ a spectral sequence E_r that converges to the $H_{\lambda\omega}^*(\mathfrak{g})$ is constructed. E_r degenerates at the first term E_1 if $-\lambda\omega$ doesn't belong to the finite set $\Omega_{\mathfrak{g}} \subset H^1(\mathfrak{g})$. $\Omega_{\mathfrak{g}}$ is defined by the collection $\{\alpha_{k+1}, \dots, \alpha_n\}$ of the closed 1-forms that have invariant sense as the weights of completely reducible representation associated to the restriction $ad|_{[\mathfrak{g}, \mathfrak{g}]}$ of adjoint representation $ad : \mathfrak{g} \rightarrow \mathfrak{g}$ (Theorem 3.3).

The main result of this article (Theorem 4.13): *the cohomology with local coefficients $H_{\lambda\omega}^*(G/\Gamma, \mathbb{C})$ of a compact solvmanifold G/Γ , where G is completely solvable Lie group is non-trivial if and only if $-\lambda[\omega] \in \tilde{\Omega}_{\mathfrak{g}}$, where $\tilde{\Omega}_{\mathfrak{g}}$ is the finite subset in $H^1(G/\Gamma, \mathbb{C})$ well defined in terms of $\mathfrak{g}$.*

The author is grateful to L. Alania for helpful discussions and attention to this work.

1. DEFORMED DERHAM COMPLEX AND MORSE-NOVIKOV THEORY

Let us consider a closed compact C^∞ -manifold M and its deRham complex $(\Lambda^*(M), d)$ of differential forms. Let ω be a closed 1-form on M and $\lambda \in \mathbb{R}$. Now one can define a new algebraic complex $(\Lambda^*(M), d_{\lambda\omega})$ with a deformed differential

$$d_{\lambda\omega} = d + \lambda\omega : \Lambda^*(M) \rightarrow \Lambda^*(M)$$

i.e. for any form $a \in \Lambda^*(M)$:

$$d_{\lambda\omega}(a) = da + \lambda\omega \wedge a.$$

Now taking $\lambda \in \mathbb{C}$ and considering the complexification $\Lambda^*(M) \otimes \mathbb{C}$ we come to the following important

Lemma 1.1 ([N3], [Pa]). *1) For a closed 1-form $d\omega = 0$ the cohomology $H_{\lambda\omega}^*(M, \mathbb{C})$ of the algebraic complex $(\Lambda^*(M) \otimes \mathbb{C}, d_{\lambda\omega})$ coincides with the cohomology $H_{\rho_{\lambda\omega}}^*(M, \mathbb{C})$ with coefficients in local system of groups $\mathbb{C}$ defined by the representation $\rho_{\lambda\omega} : \pi_1(M) \rightarrow \mathbb{C}^*$ of fundamental group defined by the formula*

$$\rho_{\lambda\omega}([\gamma]) = \exp \int_{\gamma} \lambda\omega, \quad [\gamma] \in \pi_1(M),$$

2) For any pair ω, ω' of 1-forms such that $\omega - \omega' = d\phi, \phi \in C^\infty(M)$ the cohomology $H_{\lambda\omega}^*(M, \mathbb{C})$ and $H_{\lambda\omega'}^*(M, \mathbb{C})$ are isomorphic to each other. This isomorphism can be given by the gauge transformation

$$a \rightarrow e^{\lambda\phi}a; \quad d \rightarrow e^{\lambda\phi}de^{-\lambda\phi} = d + \lambda d\phi \wedge$$

We denote corresponding Betti numbers by $b_p(\lambda, \omega)$, where $b_p(\lambda, \omega) = \dim H_{\rho_{\lambda\omega}}^*(M, \mathbb{C})$.

Remark. The representation $\rho_{\lambda\omega} : \pi_1(M) \rightarrow \mathbb{C}^*$ defines a local system of groups $\mathbb{C}$ on the manifold M in the sense of Steenrod (see [R] for details).

The cohomology $H_{\lambda\omega}^*(M, \mathbb{C})$ naturally arises in the Morse-Novikov theory: we assume now that ω is a Morse 1-form, i.e., in a neighbourhood of any point $\omega = df$, where f is a Morse function. The zeros of ω are isolated, and one can define the index of each zero. The number of zeros of ω of index p is denoted by $m_p(\omega)$.

Theorem 1.2 (A. Pazhitnov, [Pa]). *For sufficiently large real numbers λ ,*

$$m_p(\omega) \geq b_p(\lambda, \omega) \quad \forall p.$$

Theorem 1.3 (A. Pazhitnov, [Pa]). *Assume that all the periods of the form ω are commensurable. If $\text{Re}\lambda$ is sufficiently large, then $b_p(\lambda, \omega) = b_p(\omega)$, where $b_p(\omega)$ is a Novikov number.*

2. DIXMIER'S EXACT SEQUENCE OF LIE ALGEBRA COHOMOLOGY

Let $\mathfrak{g}$ be a n -dimensional Lie algebra. The dual of the Lie bracket $[\cdot, \cdot] : \Lambda^2(\mathfrak{g}) \rightarrow \mathfrak{g}$ gives a linear mapping $d_1 : \mathfrak{g}^* \rightarrow \Lambda^2(\mathfrak{g}^*)$ which extends in a standard way to a differential d of a cochain complex of the Lie algebra $\mathfrak{g}$:

$$\mathbb{K} \xrightarrow{d_0=0} \mathfrak{g}^* \xrightarrow{d_1} \Lambda^2(\mathfrak{g}^*) \xrightarrow{d_2} \Lambda^3(\mathfrak{g}^*) \xrightarrow{d_3} \dots$$

$$d(\rho \wedge \eta) = d\rho \wedge \eta + (-1)^{\deg \rho} \rho \wedge d\eta, \quad \forall \rho, \eta \in \Lambda^*(\mathfrak{g}^*).$$

Vanishing of the d^2 corresponds to the Jacobi identity.

For $d : \Lambda^q(\mathfrak{g}^*) \rightarrow \Lambda^{q+1}(\mathfrak{g}^*)$ and $f \in \Lambda^q(\mathfrak{g}^*)$ the following formula holds on:

$$(1) \quad \sum_{1 \leq i < j \leq q+1} (-1)^{i+j-1} f([X_i, X_j], X_1, \dots, \hat{X}_i, \dots, \hat{X}_j, \dots, X_{q+1}).$$

A cohomology of this complex is called the cohomology (with trivial coefficients) of the Lie algebra $\mathfrak{g}$ and is denoted by $H^*(\mathfrak{g})$.

From the definition it follows that $H^1(\mathfrak{g})$ is the dual space to $\mathfrak{g}/[\mathfrak{g}, \mathfrak{g}]$ and so

- 1) $b^1(\mathfrak{g}) = \dim H^1(\mathfrak{g}) \geq 2$ for a nilpotent Lie algebra $\mathfrak{g}$,
- 2) $b^1(\mathfrak{g}) \geq 1$ for a solvable Lie algebra $\mathfrak{g}$,
- 3) $b^1(\mathfrak{g}) = 0$ for a semi-simple Lie algebra $\mathfrak{g}$.

Now we take a Lie algebra $\mathfrak{g}$ over a field $\mathbb{K}$ with a non-trivial $H^1(\mathfrak{g})$. Let $\omega \in \mathfrak{g}^*$, $\omega \neq 0$, $d\omega = 0$ and $\lambda \in \mathbb{K}$. One can define

- 1) a new deformed differential $d_{\lambda\omega}$ in $\Lambda^*(\mathfrak{g}^*)$ by the formula

$$d_{\lambda\omega}(a) = da + \lambda\omega \wedge a.$$

- 2) an one-dimensional representation

$$\rho_{\lambda\omega} : \mathfrak{g} \rightarrow \mathbb{K}, \quad \rho_{\lambda\omega}(\xi) = \lambda\omega(\xi), \quad \xi \in \mathfrak{g}.$$

Now we recall the definition of the Lie algebra cohomology associated with a representation. Let $\mathfrak{g}$ be a Lie algebra and $\rho : \mathfrak{g} \rightarrow \mathfrak{gl}(V)$ its linear representation. We denote by $C^q(\mathfrak{g}, V)$ the space of q -linear alternating mappings of $\mathfrak{g}$ into V . Then one can consider an algebraic complex:

$$V = C^0(\mathfrak{g}, V) \xrightarrow{d_0} C^1(\mathfrak{g}, V) \xrightarrow{d_1} C^2(\mathfrak{g}, V) \xrightarrow{d_2} \dots$$

where the differential d_q is defined by:

$$(2) \quad \begin{aligned} (d_q f)(X_1, \dots, X_{q+1}) &= \sum_{i=1}^{q+1} (-1)^{i+1} \rho(X_i) (f(X_1, \dots, \hat{X}_i, \dots, X_{q+1})) + \\ &+ \sum_{1 \leq i < j \leq q+1} (-1)^{i+j-1} f([X_i, X_j], X_1, \dots, \hat{X}_i, \dots, \hat{X}_j, \dots, X_{q+1}). \end{aligned}$$

The cohomology of the complex $(C^*(\mathfrak{g}, V), d)$ is called the cohomology of the Lie algebra $\mathfrak{g}$ associated to the representation $\rho : \mathfrak{g} \rightarrow V$.

Proposition 2.1. *Let $\mathfrak{g}$ be a Lie algebra and ω is a closed 1-form. Then the complex $(\Lambda^*(\mathfrak{g}^*), d_{\lambda\omega})$ coincides with the cochain complex of the Lie algebra $\mathfrak{g}$ associated with one-dimensional representation $\rho_{\lambda\omega} : \mathfrak{g} \rightarrow \mathbb{K}$, where $\rho_{\lambda\omega}(\xi) = \lambda\omega(\xi)$, $\xi \in \mathfrak{g}$.*

The proof follows from

$$(\lambda\omega \wedge a)(X_1, \dots, X_{q+1}) = \sum_{i=1}^{q+1} (-1)^{i+1} \lambda\omega(X_i) (a(X_1, \dots, \hat{X}_i, \dots, X_{q+1})).$$

One can deduce that $H_{\lambda\omega}^0(\mathfrak{g}) = 0$ for a non-trivial $\lambda\omega$, as well as $H_{\lambda\omega}^n(\mathfrak{g}) = 0$ for an unimodular n -dimensional Lie algebra $\mathfrak{g}$.

Remark. The cohomology $H_{\lambda\omega}^*(\mathfrak{g})$ coincides with the Lie algebra cohomology $H^*(\mathfrak{g})$ with trivial coefficients if $\lambda = 0$. If $\lambda \neq 0$ the deformed differential $d_{\lambda\omega}$ is not compatible with the exterior product $\wedge$ in $\Lambda^*(\mathfrak{g})$

$$d_{\lambda\omega}(a \wedge b) = d(a \wedge b) + \lambda\omega \wedge a \wedge b \neq d_{\lambda\omega}(a) \wedge b + (-1)^{\deg a} a \wedge d_{\lambda\omega}(b)$$

and the cohomology $H_{\lambda\omega}^*(\mathfrak{g})$ has no natural multiplicative structure and therefore no Poincaré duality in the case of unimodular Lie algebra $\mathfrak{g}$. The corresponding Euler characteristic $\chi_{\lambda\omega}(\mathfrak{g})$ is still equal to zero.

Let $\omega \in \mathfrak{g}^*, \omega \neq 0, d\omega = 0$. Then $\mathfrak{b}_\omega = \{x \in \mathfrak{g}, \omega(x) = 0\}$ is an ideal of codimension 1 in $\mathfrak{g}$. One can choose an element $X \in \mathfrak{g}, \omega(X) = 1$.

Theorem 2.2 (Dixmier [D]). *There exists a long exact sequence of Lie algebra cohomology:*

$$(3) \quad \dots \xrightarrow{adX_{i-1}^* + \lambda Id} H^{i-1}(\mathfrak{b}_\omega) \xrightarrow{\omega \wedge} H_{\lambda\omega}^i(\mathfrak{g}) \xrightarrow{r_i} H^i(\mathfrak{b}_\omega) \xrightarrow{adX_i^* + \lambda Id} H^i(\mathfrak{b}_\omega) \rightarrow \dots$$

where

- 1) the homomorphism $r_i : H^i(\mathfrak{g}) \rightarrow H^i(\mathfrak{b})$ is the restriction homomorphism;
- 2) $\omega \wedge : H^{*-1}(\mathfrak{b}_\omega) \rightarrow H^*(\mathfrak{g})$ is induced by the multiplication $\omega \wedge : \Lambda^{*-1}(\mathfrak{b}_\omega^*) \rightarrow \Lambda^*(\mathfrak{g}^*)$;
- 3) the homomorphisms $adX_i^* : H^i(\mathfrak{b}_\omega) \rightarrow H^i(\mathfrak{b}_\omega)$ are induced by the derivation adX_i^* of degree zero of $\Lambda^*(\mathfrak{b}_\omega^*)$ ($adX^*(a \wedge b) = adX^*a \wedge b + a \wedge adX^*b, \forall a, b \in \Lambda^*(\mathfrak{b}_\omega^*)$) that continues a dual mapping $adX^* : \mathfrak{b}_\omega^* \rightarrow \mathfrak{b}_\omega^*$ to the $adX : \mathfrak{b}_\omega \rightarrow \mathfrak{b}_\omega$ operator. The derivation adX^* commutes with d and corresponding mapping in $H^*(\mathfrak{b}_\omega)$ we denote by the same symbol adX^* . Id is the identity operator.

Each form $f \in \Lambda^*(\mathfrak{g})$ can be decomposed as $f = \omega \wedge f' + f''$, where $f' \in \Lambda^{*-1}(\mathfrak{b}_\omega^*)$ and $f'' \in \Lambda^*(\mathfrak{b}_\omega^*)$. And one can write out a short exact sequence of algebraic complexes

$$0 \longrightarrow \Lambda^{*-1}(\mathfrak{b}_\omega^*) \xrightarrow{\omega \wedge} \Lambda^*(\mathfrak{g}^*) \longrightarrow \Lambda^*(\mathfrak{b}_\omega^*) \longrightarrow 0$$

where $\Lambda^*(\mathfrak{b}_\omega^*)$ has the standard differential d , $\Lambda^*(\mathfrak{g}^*)$ has the deformed differential $d_{\lambda\omega}$ and $\Lambda^{*-1}(\mathfrak{b}_\omega^*)$ is taken with the differential $-d$ as $d(\omega \wedge c) = d_{\lambda\omega}(\omega \wedge c) = -\omega \wedge dc$.

The short exact sequence of algebraic complexes gives us the long exact sequence of cohomology. Everything is clear with the homomorphisms r_i and $\omega \wedge$. As for the homomorphism $H^q(\mathfrak{b}_\omega) \rightarrow H^q(\mathfrak{b}_\omega)$ we can remark the following.

First of all let us introduce a new mapping

$$\Lambda^*(\mathfrak{g}^*) \rightarrow \Lambda^{*-1}(\mathfrak{b}_\omega^*), f \in \Lambda^*(\mathfrak{g}^*) \rightarrow f_X \in \Lambda^{*-1}(\mathfrak{b}_\omega^*),$$

where $f_X(X_1, \dots, X_q) = f(X, X_1, \dots, X_q)$. So $f_X = f'$ if $f = \omega \wedge f' + f''$.

Then an obvious formula holds on:

$$\begin{aligned} (4) \quad (df)_X(X_1, \dots, X_{q+1}) &= \sum_{1 \leq j \leq q+1} (-1)^j f(adX(X_j), X_1, \dots, \hat{X}_j, \dots, X_{q+1}) + \\ &+ \sum_{1 \leq i < j \leq q+1} (-1)^{i+j} f([X_i, X_j], X, X_1, \dots, \hat{X}_i, \dots, \hat{X}_j, \dots, X_{q+1}) = \\ &= (adX_q^*(f) + d(f_X))(X_1, \dots, X_{q+1}). \end{aligned}$$

Hence the homomorphism $H^q(\mathfrak{b}_\omega) \rightarrow H^q(\mathfrak{b}_\omega)$, $[f] \rightarrow [f' + \lambda f]$ of long exact sequence in cohomology coincides with homomorphism induced by $adX^* + \lambda Id$.

Corollary 2.3. *Let $\mathfrak{g}$ be a n -dimensional Lie algebra and $\omega \in \mathfrak{g}^*$, $\omega \neq 0$, $d\omega = 0$ and $Spec^k(\omega)$ be the set of eigenvalues of operator $adX_k^* : H^k(\mathfrak{b}_\omega) \rightarrow H^k(\mathfrak{b}_\omega)$, then*

1) *the cohomology $H_{\lambda\omega}^*(\mathfrak{g})$ is non-trivial if and only if*

$$-\lambda \in \cup_{k=1}^n Spec^k(\omega);$$

2) *the i -th Betti number $b_{\lambda\omega}^i(\mathfrak{g}) = \dim H_{\lambda\omega}^i(\mathfrak{g})$ can be expressed in a following way:*

$$(5) \quad b_{\lambda\omega}^i(\mathfrak{g}) = k_{\lambda\omega}^i + k_{\lambda\omega}^{i-1},$$

where by $k_{\lambda\omega}^i$ we denote the dimension of the kernel of $adX_i^* + \lambda Id : H^i(\mathfrak{b}_\omega) \rightarrow H^i(\mathfrak{b}_\omega)$.

Example 2.4. Let $\mathfrak{g}$ be a Lie algebra defined by the basis $X, e_1, e_2, \dots, e_n$ and commuting relations (trivial ones are omitted):

$$[X, e_1] = e_1, [X, e_2] = e_2, \dots, [X, e_n] = e_n.$$

Thus $\mathfrak{g}$ is a semidirect sum of $\mathbb{K}$ and abelian $\mathbb{K}^n$ defined by the operator adX with the identity matrix E in the basis $e_1, e_2, \dots, e_n$ of $\mathbb{K}^n$. We take $\omega, \omega_1, \dots, \omega_n$ as the corresponding dual basis in $\mathfrak{g}$. In particular

$$\omega(X) = 1, \omega(e_i) = 0, i = 1, \dots, n.$$

$d\omega = 0$ and $adX^*(\omega_{i_1} \wedge \dots \wedge \omega_{i_p}) = p\omega_{i_1} \wedge \dots \wedge \omega_{i_p}$. Hence $Spec^p(\omega) = \{p\}$ and for $\lambda = p$ we have only two non-trivial Betti numbers:

$$b_{p\omega}^{p-1}(\mathfrak{g}) = b_{p\omega}^p(\mathfrak{g}) = \binom{n}{p}.$$

Corollary 2.5 (Dixmier, [D]). *Let $\mathfrak{g}$ be a nilpotent Lie algebra. The cohomology $H_{\lambda\omega}^*(\mathfrak{g})$ is trivial for all non-trivial $\lambda\omega$ and coincides with the Lie algebra cohomology $H^*(\mathfrak{g})$ in trivial $\lambda\omega = 0$ case.*

The operator adX is nilpotent and therefore the same is $adX_i^* : H^i(\mathfrak{b}_\omega) \rightarrow H^i(\mathfrak{b}_\omega)$. Hence $adX_i^* + \lambda Id$ is non-degenerate operator for all $\lambda \neq 0$. We obtain the proof of Theorem 1 from [Al] as the corollary of Dixmier's theorem [D] for cohomology of nilpotent Lie algebras.

Remark. We represented in this article only a version of Dixmier's exact sequence adapted to our special case of 1-dimensional Lie algebra representation (see [D] for all details), the last thing that we want to recall is Dixmier's estimate for Betti numbers $\dim H^p(\mathfrak{g})$ of a nilpotent Lie algebra $\mathfrak{g}$.

Corollary 2.6 ([D]). *Let $\mathfrak{g}$ be a nilpotent Lie algebra. Then*

$$\dim H^p(\mathfrak{g}) \geq 2, \quad p = 1, \dots, n-1.$$

It follows from the Corollary 2.3: we have $\lambda\omega = 0$ and $\dim H^p(\mathfrak{g}) = k^p + k^{p-1}$, where $k^p \geq 1, k^{p-1} \geq 1$ are the dimensions of the kernels of nilpotent operators adX_p^*, adX_{p-1}^* in the spaces $H^p(\mathfrak{b}_\omega), H^{p-1}(\mathfrak{b}_\omega)$ with $\dim H^p(\mathfrak{b}_\omega) \geq 2, \dim H^{p-1}(\mathfrak{b}_\omega) \geq 2$ by inductive assumption.

3. COHOMOLOGY OF SOLVABLE LIE ALGEBRAS

Definition 3.1. A real solvable Lie algebra $\mathfrak{g}$ is called completely solvable if $ad(X) : \mathfrak{g} \rightarrow \mathfrak{g}$ has only real eigenvalues $\forall X \in \mathfrak{g}$.

Lemma 3.2. *Let $\mathfrak{g}$ be a n -dimensional solvable over $\mathbb{C}$ (or real completely solvable Lie algebra) and $b^1(\mathfrak{g}) = \dim H^1(\mathfrak{g}) = k$. Then exists a basis $\omega_1, \dots, \omega_n$ in $\mathfrak{g}^*$ such that*

$$(6) \quad d\omega_1 = \dots = d\omega_k = 0,$$

$$d\omega_{k+1} = \alpha_{k+1} \wedge \omega_{k+1} + P_{k+1}(\omega_1, \dots, \omega_k),$$

$$\dots,$$

$$d\omega_n = \alpha_n \wedge \omega_n + P_n(\omega_1, \dots, \omega_{n-1}),$$

where $\alpha_{k+1}, \dots, \alpha_n$ are closed 1-forms, that are the weights of completely reducible representation associated to $ad|_{[\mathfrak{g}, \mathfrak{g}]}$ and $P_i(\omega_1, \dots, \omega_{i-1}) \in \Lambda^2(\omega_1, \dots, \omega_{i-1})$.

For the proof we apply Lie's theorem to the adjoint representation ad restricted to the commutant $[\mathfrak{g}, \mathfrak{g}]$. Namely we can choose a basis $e_{k+1}, \dots, e_n$ such that the subspaces $Span(e_i, \dots, e_n), i = k+1, \dots, n$ are invariant with respect to the representation $ad|_{[\mathfrak{g}, \mathfrak{g}]}$. Then we add $e_1, \dots, e_k$ in a way that $e_1, \dots, e_n$ form the basis of $\mathfrak{g}$. For the dual forms $\omega_1, \dots, \omega_n$ in $\mathfrak{g}^*$ we have formulas (6).

Remark. One can consider the free d -algebra $(\Lambda^*(\omega_1, \dots, \omega_n), d)$ as a kind of minimal model of the cochain complex $\Lambda^*(\mathfrak{g}^*)$ because the mapping $\Lambda^*(\omega_1, \dots, \omega_n) \rightarrow \Lambda^*(\mathfrak{g}^*)$ induces the isomorphism in cohomology.

Now we start with a solvable Lie algebra over $\mathbb{C}$ (or real completely solvable) and take a basis $\omega_1, \dots, \omega_n$ constructed in Lemma 3.2. Let us denote by Λ^* the exterior subalgebra in $\Lambda^*(\omega_1, \dots, \omega_n)$ generated by $\omega_1, \dots, \omega_k$. One can define a filtration F of the cochain complex $\Lambda^*(\mathfrak{g}^*)$:

$$0 \subset \Lambda^* \subset F^{k+1} \subset F^{k+2} \subset F^{k+1, k+2} \subset F^{k+3} \subset F^{k+1, k+3} \subset \dots \subset F^{k, \dots, n} = \Lambda^*(\mathfrak{g}^*)$$

where the system of subspaces $\{F^{j_1, \dots, j_p}, k < j_1 < \dots < j_p \leq n\}$ is defined by the following conditions:

A subspace $F^{j_1, \dots, j_p}$ is spanned by monomials $a = \omega_{l_1} \wedge \dots \wedge \omega_{l_q}$ with $l_1 < \dots < l_q$ such that

$$\begin{aligned} & l_q < j_p; \text{ or } l_q = j_p, l_{q-1} < j_{p-1}; \\ & \text{or } l_q = j_p, l_{q-1} = j_{p-1}, l_{q-2} < j_{p-2}; \text{ or } \dots; \\ & \text{or } l_q = j_p, \dots, l_{q-p+1} = j_1, l_{q-p} \leq k. \end{aligned}$$

Thus for example:

$$\begin{aligned} (7) \quad F^{k+1} &= \Lambda^* \oplus \Lambda^* \wedge \omega_{k+1}, F^{k+2} = \Lambda^* \oplus \Lambda^* \wedge \omega_{k+1} \oplus \Lambda^* \wedge \omega_{k+2}, \\ F^{k+1, k+2} &= \Lambda^* \oplus \Lambda^* \wedge \omega_{k+1} \oplus \Lambda^* \wedge \omega_{k+2} \oplus \Lambda^* \wedge \omega_{k+1} \wedge \omega_{k+2}, \\ F^{k+3} &= \Lambda^* \oplus \Lambda^* \wedge \omega_{k+1} \oplus \Lambda^* \wedge \omega_{k+2} \oplus \Lambda^* \wedge \omega_{k+1} \wedge \omega_{k+2} \oplus \Lambda^* \wedge \omega_{k+3}, \dots \end{aligned}$$

The filtration F is compatible with differential $d + \lambda\omega$ and one can consider the corresponding spectral sequence E_r . To obtain its first term E_1 one have to calculate the cohomology of complexes $(\Lambda^* \wedge \omega_{j_1} \wedge \dots \wedge \omega_{j_p}, d_0)$:

$$d_0(\tilde{a} \wedge \omega_{j_1} \wedge \dots \wedge \omega_{j_p}) = (\alpha_{j_1} + \dots + \alpha_{j_p} + \lambda\omega)\tilde{a} \wedge \omega_{j_1} \wedge \dots \wedge \omega_{j_p}.$$

The cohomology $H^*(\Lambda^* \wedge \omega_{j_1} \wedge \dots \wedge \omega_{j_p}, d_0)$ coincides with the cohomology of $(\Lambda^*, \tilde{d})$ where differential $\tilde{d}$ acts as exterior multiplication by 1-form $\alpha_{j_1} + \dots + \alpha_{j_p} + \lambda\omega$. Hence $H^*(\Lambda^* \wedge \omega_{j_1} \wedge \dots \wedge \omega_{j_p}, d_0)$ is trivial if $\alpha_{j_1} + \dots + \alpha_{j_p} + \lambda\omega \neq 0$. So taking $\lambda\omega$ such that

$$\alpha_{j_1} + \dots + \alpha_{j_p} + \lambda\omega \neq 0 \quad \forall j_1 < \dots < j_p, p = k+1, \dots, n$$

one can see that the spectral sequence E_r degenerates at the first term E_1 . Taking the complexification of real solvable Lie algebra $\mathfrak{g}$ we come to the following

Theorem 3.3. *Let $\mathfrak{g}$ be a solvable Lie algebra, $\dim \mathfrak{g} = n$ and $\{\alpha_{k+1}, \dots, \alpha_n\}$ is the collection of the weights of completely reducible representation associated to $\text{ad}|_{[\mathfrak{g}, \mathfrak{g}]}$. Let $\Omega_{\mathfrak{g}}$ denote the set of all p -sums $\alpha_{i_1} + \dots + \alpha_{i_p}$, $k+1 \leq i_1 < \dots < i_p \leq n$, $p = 1, \dots, n$. Then exists a spectral sequence E_r that converges to the cohomology $H_{\lambda\omega}^*(\mathfrak{g})$ and its first term E_1 degenerates if $-\lambda\omega \notin \Omega_{\mathfrak{g}}$.*

The set $\Omega_{\mathfrak{g}}$ is defined by $\lambda\omega$ such that the term E_1 is non-trivial, but generally E_1 doesn't coincide with E_{∞} . So we have to introduce $\tilde{\Omega}_{\mathfrak{g}} \subset \Omega_{\mathfrak{g}}$ such that $E_{\infty} \neq 0$ if and only if $-\lambda\omega \in \{0\} \cup \tilde{\Omega}_{\mathfrak{g}}$.

Corollary 3.4. *Let $\mathfrak{g}$ be a solvable Lie algebra. Then $H_{\rho_{\lambda\omega}}^*(\mathfrak{g})$ is non-trivial if and only if $-\lambda\omega \in \{0\} \cup \tilde{\Omega}_{\mathfrak{g}}$ – the finite subset in $H^1(\mathfrak{g})$.*

4. COHOMOLOGY OF SOLVMANIFOLDS.

Definition 4.1. A solvmanifold (nilmanifold) M is a compact homogeneous space of the form G/Γ , where G is a simply connected solvable (nilpotent) Lie group and Γ is a lattice in G .

Let $\mathfrak{g}$ denote a Lie algebra of G . Recall that G is solvable if and only if $\mathfrak{g}$ is solvable Lie algebra, the last condition is equivalent to nilpotency of derived algebra $[\mathfrak{g}, \mathfrak{g}]$.

We start with examples of nilmanifolds.

Example 4.2. A n -dimensional torus $T^n = \mathbb{R}^n / \mathbb{Z}^n$.

Example 4.3. The Heisenberg manifold $M_3 = \mathcal{H}_3/\Gamma_3$, where $\mathcal{H}_3$ is the group of all matrices of the form

$$\begin{pmatrix} 1 & x & z \\ 0 & 1 & y \\ 0 & 0 & 1 \end{pmatrix}, \quad x, y, z \in \mathbb{R},$$

and a lattice Γ_3 is a subgroup of matrices with $x, y, z \in \mathbb{Z}$.

Theorem 4.4 (A.I. Malcev [Mal]). *Let G be a simply connected nilpotent Lie group with a tangent Lie algebra $\mathfrak{g}$. Then G has a co-compact lattice Γ (i.e. G/Γ is a compact space) if and only if there exists a basis $e_1, e_2, \dots, e_n$ in $\mathfrak{g}$ such that the constants $\{c_{ij}^k\}$ of Lie structure $[e_i, e_j] = c_{ij}^k e_k$ are all rational numbers.*

This theorem gives us a practical tool for construction of nilmanifolds: let $\mathfrak{g}$ be a nilpotent Lie algebra defined by its basis $e_1, e_2, \dots, e_n$ and commuting relations $[e_i, e_j] = c_{ij}^k e_k$, where all numbers $c_{ij}^k \in \mathbb{Q}$. Now one can define a group structure $*$ in the vector space $\mathfrak{g}$ using the Campbell-Hausdorff formula. The nilpotent group $G = (\mathfrak{g}, *)$ has a co-compact lattice Γ (a subgroup generated by basic elements $e_1, e_2, \dots, e_n$) and one can consider corresponding nilmanifold G/Γ .

Example 4.5. Let $\mathcal{V}_n$ be a nilpotent Lie algebra with a basis $e_1, e_2, \dots, e_n$ and a Lie bracket:

$$[e_i, e_j] = \begin{cases} (j-i)e_{i+j}, & i+j \leq n; \\ 0, & i+j > n. \end{cases}$$

Cohomology of the corresponding family of nilmanifolds M_n was studied in [Al] and [Mill].

The situation with non-nilpotent solvable Lie groups is much more difficult: the crucial point is the problem of existence of cocompact lattice (see [R] for details). For example, if a solvable Lie group G admits a cocompact lattice Γ then the corresponding Lie algebra $\mathfrak{g}$ is unimodular, hence $\alpha_{k+1} + \dots + \alpha_n = 0$. The condition of unimodularity of $\mathfrak{g}$ is not sufficient. See [R] for general information in this domain.

Example 4.6. Let us consider a semidirect product $G_0 = \mathbb{R} \ltimes \mathbb{R}^2$ where $\mathbb{R}$ acts on $\mathbb{R}^2$ via

$$t \rightarrow \phi(t) = \begin{pmatrix} a^t & 0 \\ 0 & a^{-t} \end{pmatrix},$$

where $a + a^{-1} = n \in \mathbb{N}, a \neq 1, a > 0$. Then

$$\phi(1) = C^{-1} \begin{pmatrix} 0 & 1 \\ -1 & n \end{pmatrix} C,$$

for some matrix $C \in GL(2, \mathbb{R})$. Then exists a lattice $L \subset \mathbb{R}^2$ invariant with respect to $\phi(1)$. $\mathbb{Z}$ acts on L via $\phi(1)^m$ and we define a lattice $\Gamma \subset G$ as $\mathbb{Z} \ltimes_{\phi(1)} L$. The lattices corresponding to different values of n are, generally speaking, non-isomorphic. So corresponding solvmanifolds are non diffeomorphic to. But we study the cohomology $H_\rho^*(G/\Gamma, \mathbb{C})$ over $\mathbb{C}$ and as we will see it doesn't depend on the choice of $\Gamma \subset G$.

The solvmanifold from the previous example is a fibre bundle over S^1 with T^2 as fibre. It can be generalized by the following

Theorem 4.7 (G.D. Mostow [Mos1]). *Any compact solvmanifold is a bundle with toroid as base space and nilmanifold as fibre.*

Definition 4.8. A solvable Lie group G is called completely solvable if its tangent Lie algebra $\mathfrak{g}$ is completely solvable.

One can identify deRham complex $\Lambda^*(G/\Gamma)$ with subcomplex $\Lambda_\Gamma^*(G) \subset \Lambda^*(G)$ of left-invariant forms on G with respect to the action of the lattice Γ . $\Lambda_\Gamma^*(G)$ contains the subcomplex $\Lambda_G^*(G)$ of left-invariant forms with respect to the whole action of G . $\Lambda_G^*(G)$ is naturally isomorphic to the Lie algebra cochain complex $\Lambda^*(\mathfrak{g})$. Let us consider the corresponding inclusion

$$\psi : \Lambda^*(\mathfrak{g}) \rightarrow \Lambda^*(G/\Gamma).$$

Theorem 4.9 (Hattori [H]). *Let G/Γ be a compact solvmanifold, where G is a completely solvable Lie group, then the inclusion $\psi : \Lambda^*(\mathfrak{g}) \rightarrow \Lambda^*(G/\Gamma)$ induces the isomorphism $\psi^* : H^*(\mathfrak{g}) \rightarrow H^*(G/\Gamma, \mathbb{R})$ in cohomology.*

Remark. In fact Hattori's theorem is the generalization of the theorem proved by Nomizu [Nz] for nilmanifolds. For an arbitrary solvmanifold G/Γ the mapping ψ^* is not isomorphism but it is an inclusion (see [R]).

So every class $[\omega] \in H^1(G/\Gamma, \mathbb{R})$ can be represented by the left-invariant (with respect to the action of G) 1-form ω . By means of ω one can define a one-dimensional representation $\rho_{\lambda\omega} : G \rightarrow \mathbb{C}^*$:

$$\rho_{\lambda\omega}(g) = \exp \int_{\gamma(e,g)} \lambda\omega,$$

where $\gamma(e, g)$ is a path connecting the identity e with $g \in G$ (let us recall that G is a simply connected). As ω is the left invariant 1-form then

$$\int_{\gamma(e, g_1 g_2)} \lambda\omega = \int_{\gamma(e, g_1)} \lambda\omega + \int_{\gamma(g_1, g_1 g_2)} \lambda\omega = \int_{\gamma(e, g_1)} \lambda\omega + \int_{g_1^{-1} \gamma(e, g_2)} \lambda\omega$$

holds on and $\rho_{\lambda\omega}(g_1 g_2) = \rho_{\lambda\omega}(g_1) \rho_{\lambda\omega}(g_2)$. $\rho_{\lambda\omega}$ induces the representation of corresponding Lie algebra $\mathfrak{g}$ (we denote it by the same symbol): $\rho_{\lambda\omega}(X) = \lambda\omega(X)$.

In this situation it's possible to make some generalizations using Mostow's theorem. Namely following [R] we give

Definition 4.10. Let G be a simply-connected Lie group and $\Gamma \subset G$ a lattice. Let ρ be a finite dimensional representation of G on a complex vector space F . Let Ad denote the adjoint representation of G on its Lie algebra $\mathfrak{g}$ as well as the complexification $\mathfrak{g}_{\mathbb{C}}$ of $\mathfrak{g}$. We will say that the representation ρ is Γ -supported if $\rho(\Gamma)$ and $\rho(G)$ have the same Zariski closure in $Aut_{\mathbb{C}}(F)$. The representation ρ is Γ -admissible if $\rho \oplus Ad$ (on $F \oplus \mathfrak{g}_{\mathbb{C}}$) is Γ -supported.

Theorem 4.11 (Mostow, Theorem 7.26 in [R]). *Let G be a simply-connected solvable Lie group and a $\Gamma \subset G$ a lattice. Let ρ be a finite dimensional Γ -admissible representation in a complex vector space F .*

Then the inclusion $\psi : \Lambda^*(\mathfrak{g}, F) \rightarrow \Lambda^*(G/\Gamma, F)$ of complexes of differential forms with values in F induces the isomorphism $\psi^* : H_\rho^*(\mathfrak{g}, F) \rightarrow H_\rho^*(G/\Gamma, F)$ in cohomology where ρ is used also to denote the representation of the Lie algebra $\rho : \mathfrak{g} \rightarrow F$ induced by $\rho : G \rightarrow Aut_{\mathbb{C}}(F)$.

But we'll not discuss the details of Mostow's theorem and possible generalizations we restrict ourselves to the case of completely solvable Lie group G . Namely we'll prove by means of Hattori's theorem the following important

Corollary 4.12. *Let G/Γ be a compact solvmanifold, G has a completely solvable Lie group and ω is a closed 1-form on G/Γ . The cohomology $H_{\lambda\omega}^*(G/\Gamma, \mathbb{C})$ is isomorphic to the Lie algebra cohomology $H_{\lambda\omega'}^*(\mathfrak{g})$ where $\omega' \in \mathfrak{g}^*$ is the left-invariant 1-form that represents the class $[\omega] \in H^1(G/\Gamma, \mathbb{R})$.*

The Corollary 4.12 together with Corollary 3.4 gives us

Theorem 4.13. *Let G/Γ be a compact solvmanifold, G is a completely solvable Lie group and ω is a closed 1-form on G/Γ . The cohomology $H_{\lambda\omega}^*(G/\Gamma, \mathbb{C})$ is non-trivial if and only if $-\lambda[\omega] \in \{0\} \cup \tilde{\Omega}_{\mathfrak{g}}$ – the finite subset in $H^1(G/\Gamma, \mathbb{R})$ well-defined in terms of the corresponding Lie algebra $\mathfrak{g}$.*

Corollary 4.14 ([Al]). *Let G/Γ be a compact nilmanifold. The cohomology $H_{\lambda\omega}^*(G/\Gamma, \mathbb{C}) = 0$ if and only if $\lambda\omega \neq 0$.*

Let us consider a 3-dimensional solvmanifold $M = G_0/\Gamma$ defined in the Example 4.6. The corresponding Lie algebra $\mathfrak{g}_0$ is isomorphic to the Lie algebra defined by its basis e_1, e_2, e_3 and following non-trivial brackets:

$$[e_1, e_2] = e_2, [e_1, e_3] = -e_3.$$

We take a dual basis $\omega_1, \omega_2, \omega_3$ in $\mathfrak{g}^*$. Then

$$d\omega_1 = 0, d\omega_2 = \omega_1 \wedge \omega_2, d\omega_3 = -\omega_1 \wedge \omega_3.$$

So $\alpha_2 = \omega_1$ and $\alpha_3 = -\omega_1$. It is easy to see that $H_{\lambda\omega}^*(\mathfrak{g}_0)$ is non-trivial if and only if $\lambda\omega = 0, \pm\omega_1$. For corresponding Betti numbers $b_{\lambda\omega}^p(M) = \dim H_{\lambda\omega}^p(G_0/\Gamma) = \dim H_{\lambda\omega}^p(\mathfrak{g}_0)$ of the solvmanifold $M = G_0/\Gamma$ we have:

$$b_{\pm\omega_1}^0(M) = 0, b_{\pm\omega_1}^1(M) = b_{\pm\omega_1}^2(M) = 1, b_{\pm\omega_1}^3(M) = 0;$$

$$b^0(M) = b^1(M) = b^2(M) = b^3(M) = 1.$$

REFERENCES

- [Al] L. Alania, *Cohomology with local system of certain nilmanifolds*, Russian Math. Surveys **54**:5 (1999), 1019–1020.
- [D] J. Dixmier, *Cohomologie des algebres de Lie nilpotentes*, Acta Sci. Math. Szeged **16** (1955), 246–250.
- [H] A. Hattori, *Spectral sequence in the deRham cohomology of fibre bundles*, J. Fac. Sci. Univ. Tokyo, Sect. 1, **8**:4, (1960), pp. 289–331.
- [Mal] A. Malcev, *On a class of homogeneous spaces*, Amer. Math. Soc. Transl. (1) **9** (1962), 276–307.
- [Mill] D.V. Millionschikov, *Cohomology of nilmanifolds and Gontcharova's theorem*, in "Global Differential geometry: The Mathematical Legacy of Alfred Gray", M. Fernandez and J. Wolf ed., AMS CONM 288 (2001), 381–385.
- [Mos1] G.D. Mostow, *Factor spaces of solvable groups*, Ann. of Math. **60** (1954), 1–27.
- [Mos2] G.D. Mostow, *Cohomology of topological groups and solvmanifolds*, Ann. of Math. **73** (1961), 20–48.
- [Nz] K. Nomizu, *On the cohomology of homogeneous spaces of nilpotent Lie groups*, Ann. of Math. **59** (1954), 531–538.
- [N1] S.P. Novikov, *Multivalued functions and functionals. An analogue of the Morse theory*, Soviet Math. Dokl. **24** (1981), 222–226.
- [N2] S.P. Novikov, *The hamiltonian formalism and a many-valued analogue of Morse theory*, Russian Math. Surveys **37**:5 (1982), 1–56.
- [N3] S.P. Novikov, *Bloch homology. Critical points of functions and closed 1-forms*, Soviet Math. Dokl. **33**:5 (1986), 551–555.

- [N4] S.P. Novikov, *On the exotic De-Rham cohomology. Perturbation theory as a spectral sequence*, arXiv:math-ph/0201019.
- [Pa] A.V. Pazhitnov, *An analytic proof of the real part of Novikov's inequalities*, Soviet Math. Dokl. **35** (1987), 1–2.
- [R] M.S. Raghunathan, *Discrete Subgroups of Lie Groups*, Springer-Verlag, 1972.
- [W] E. Witten, *Supersymmetry and Morse theory*, J. Differential Geom. **17** (1982), 661–692.

DEPARTMENT OF MATHEMATICS AND MECHANICS, MOSCOW STATE UNIVERSITY, 119899 MOSCOW, RUSSIA

Current address: Université Louis Pasteur, UFR de Mathématique et d'Informatique, 7 rue René Descartes - 67084 Strasbourg Cedex (France)

E-mail address: `million@mech.math.msu.su`